

Some Characterizations of Cylindrical Helices in $\mathbf{E}^n$

Ahmad T. Ali

Mathematics Department
Faculty of Science, Al-Azhar University
Nasr City, 11448, Cairo, Egypt
email: atali71@yahoo.com

Rafael López*

Departamento de Geometría y Topología
Universidad de Granada
18071 Granada, Spain
email: rcamino@ugr.es

January 21, 2009

Abstract

We consider a unit speed curve α in Euclidean n -dimensional space $\mathbf{E}^n$ and denote the Frenet frame by $\{\mathbf{V}_1, \dots, \mathbf{V}_n\}$. We say that α is a cylindrical helix if its tangent vector $\mathbf{V}_1$ makes a constant angle with a fixed direction U . In this work we give different characterizations of such curves in terms of their curvatures.

MSC: 53C40, 53C50

Keywords: Euclidean n -space; Frenet equations; Cylindrical helices curves.

*Partially supported by MEC-FEDER grant no. MTM2007-61775 and Junta de Andalucía grant no. P06-FQM-01642.

1 Introduction and statement of results

An helix in Euclidean 3-space $\mathbf{E}^3$ is a curve where the tangent lines make a constant angle with a fixed direction. An helix curve is characterized by the fact that the ratio κ/τ is constant along the curve, where κ and τ are the curvature and the torsion of α , respectively. Helices are well known curves in classical differential geometry of space curves [4] and we refer to the reader for recent works on this type of curves [2, 7]. Recently, Magden [3] have introduced the concept of cylindrical helix in Euclidean 4-space $\mathbf{E}^4$ saying that the tangent lines makes a constant angle with a fixed directions. He characterizes a cylindrical helix in $\mathbf{E}^4$ if and only if the function

$$\left(\frac{\kappa_1}{\kappa_2}\right)^2 + \left(\frac{1}{\kappa_3}\left(\frac{\kappa_1}{\kappa_2}\right)'\right)^2 \quad (1)$$

is constant along the curve, where κ_3 and κ_4 are the third and the fourth curvature of the the curve. See also [5].

In this work we consider the generalization of the concept of general helices in Euclidean n-space $\mathbf{E}^n$. Let $\alpha : I \subset \mathbb{R} \rightarrow \mathbf{E}^n$ be an arbitrary curve in $\mathbf{E}^n$. Recall that the curve α is said to be of unit speed (or parameterized by arc-length function s) if $\langle \alpha'(s), \alpha'(s) \rangle = 1$, where $\langle \cdot, \cdot \rangle$ is the standard scalar product in Euclidean space $\mathbf{E}^n$ given by

$$\langle X, Y \rangle = \sum_{i=1}^n x_i y_i,$$

for each $X = (x_1, \dots, x_n)$, $Y = (y_1, \dots, y_n) \in \mathbf{E}^n$.

Let $\{\mathbf{V}_1(s), \dots, \mathbf{V}_n(s)\}$ be the moving frame along α , where the vectors $\mathbf{V}_i$ are mutually orthogonal vectors satisfying $\langle \mathbf{V}_i, \mathbf{V}_i \rangle = 1$. The Frenet equations for α are given by ([2])

$$\begin{bmatrix} \mathbf{V}'_1 \\ \mathbf{V}'_2 \\ \mathbf{V}'_3 \\ \vdots \\ \mathbf{V}'_{n-1} \\ \mathbf{V}'_n \end{bmatrix} = \begin{bmatrix} 0 & \kappa_1 & 0 & 0 & \cdots & 0 & 0 \\ -\kappa_1 & 0 & \kappa_2 & 0 & \cdots & 0 & 0 \\ 0 & -\kappa_2 & 0 & \kappa_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 0 & \kappa_{n-1} \\ 0 & 0 & 0 & 0 & \cdots & -\kappa_{n-1} & 0 \end{bmatrix} \begin{bmatrix} \mathbf{V}_1 \\ \mathbf{V}_2 \\ \mathbf{V}_3 \\ \vdots \\ \mathbf{V}_{n-1} \\ \mathbf{V}_n \end{bmatrix}. \quad (2)$$

Recall the functions $\kappa_i(s)$ are called the i-th curvatures of α . If $\kappa_{n-1}(s) = 0$ for any $s \in I$, then $\mathbf{V}_n(s)$ is a constant vector V and the curve α lies in a $(n-1)$ -dimensional affine subspace orthogonal to V , which is isometric to the Euclidean $(n-1)$ -space

$\mathbf{E}^{n-1}$. We will assume throughout this work that all the curvatures satisfy $\kappa_i(s) \neq 0$ for any $s \in I$, $1 \leq i \leq n-1$.

Definition 1.1. A unit speed curve $\alpha : I \rightarrow \mathbf{E}^n$ is called *cylindrical helix* if its tangent vector $\mathbf{V}_1$ makes a constant angle with a fixed direction U .

Our main result in this work is the following characterization of cylindrical helices in Euclidean n -space $\mathbf{E}^n$.

Theorem 1.2. Let $\alpha : I \rightarrow \mathbf{E}^n$ be a unit speed curve in $\mathbf{E}^n$. Define the functions

$$G_1 = 1, \quad G_2 = 0, \quad G_i = \frac{1}{\kappa_{i-1}} \left[\kappa_{i-2} G_{i-2} + G'_{i-1} \right], \quad 3 \leq i \leq n. \quad (3)$$

Then α is a cylindrical helix if and only if the function

$$\sum_{i=3}^n G_i^2 = C \quad (4)$$

is constant. Moreover, the constant $C = \tan^2 \theta$, being θ the angle that makes $\mathbf{V}_1$ with the fixed direction U that determines α .

This theorem generalizes in arbitrary dimensions what happens for $n = 3$ and $n = 4$, namely: if $n = 3$, (4) writes $G_3^2 = \kappa_1/\kappa_2 = \kappa/\tau$ and for $n = 4$, (4) agrees with (1).

2 Proof of Theorem 1.2

Let α be a unit speed curve in $\mathbf{E}^n$. Assume that α is a cylindrical helix curve. Let U be the direction with which $\mathbf{V}_1$ makes a constant angle θ and, without loss of generality, we suppose that $\langle U, U \rangle = 1$. Consider the differentiable functions a_i , $1 \leq i \leq n$,

$$U = \sum_{i=1}^n a_i(s) \mathbf{V}_i(s), \quad s \in I, \quad (5)$$

that is,

$$a_i = \langle \mathbf{V}_i, U \rangle, \quad 1 \leq i \leq n.$$

Then the function $a_1(s) = \langle \mathbf{V}_1(s), U \rangle$ is constant, and it agrees with $\cos \theta$:

$$a_1(s) = \langle \mathbf{V}_1, U \rangle = \cos \theta \quad (6)$$

for any s . By differentiation (6) with respect to s and using the Frenet formula (2) we have

$$a'_1(s) = \kappa_1 \langle \mathbf{V}_2, U \rangle = \kappa_1 a_2 = 0.$$

Then $a_2 = 0$ and therefore U lies in the subspace $Sp(\mathbf{V}_1, \mathbf{V}_3, \dots, \mathbf{V}_n)$. Because the vector field U is constant, a differentiation in (5) together (2) gives the following ordinary differential equation system

$$\left. \begin{array}{lcl} \kappa_1 a_1 - \kappa_2 a_3 & = & 0 \\ a'_3 - \kappa_3 a_4 & = & 0 \\ a'_4 + \kappa_3 a_3 - \kappa_4 a_5 & = & 0 \\ \vdots & & \\ a'_{n-1} + \kappa_{n-2} a_{n-2} - \kappa_{n-1} a_n & = & 0 \\ a'_n + \kappa_{n-1} a_{n-1} & = & 0 \end{array} \right\} \quad (7)$$

Define the functions $G_i = G_i(s)$ as follows

$$a_i(s) = G_i(s) a_1, \quad 3 \leq i \leq n. \quad (8)$$

We point out that $a_1 \neq 0$: on the contrary, (7) gives $a_i = 0$, for $3 \leq i \leq n$ and so, $U = 0$: contradiction. The first $(n-2)$ -equations in (7) lead to

$$\left. \begin{array}{l} G_3 = \frac{\kappa_1}{\kappa_2} \\ G_4 = \frac{1}{\kappa_3} G'_3 \\ G_5 = \frac{1}{\kappa_4} [\kappa_3 G_3 + G'_4] \\ \vdots \\ G_{n-1} = \frac{1}{\kappa_{n-2}} [\kappa_{n-3} G_{n-3} + G'_{n-2}] \\ G_n = \frac{1}{\kappa_{n-1}} [\kappa_{n-2} G_{n-2} + G'_{n-1}] \end{array} \right\} \quad (9)$$

The last equation of (7) leads to the following condition;

$$G'_n + \kappa_{n-1} G_{n-1} = 0. \quad (10)$$

We do the change of variables:

$$t(s) = \int^s \kappa_{n-1}(u) du, \quad \frac{dt}{ds} = \kappa_{n-1}(s).$$

In particular, and from the last equation of (9), we have

$$G'_{n-1}(t) = G_n(t) - \left(\frac{\kappa_{n-2}(t)}{\kappa_{n-1}(t)} \right) G_{n-2}(t).$$

As a consequence, if α is a cylindrical helix, substituting the equation (10) in the last equation yields

$$G''_n(t) + G_n(t) = \frac{\kappa_{n-2}(t)G_{n-2}(t)}{\kappa_{n-1}(t)}.$$

The general solution of this equation is

$$G_n(t) = \left(A - \int \frac{\kappa_{n-2}(t)G_{n-2}(t)}{\kappa_{n-1}(t)} \sin t \, dt \right) \cos t + \left(B + \int \frac{\kappa_{n-2}(t)G_{n-2}(t)}{\kappa_{n-1}(t)} \cos t \, dt \right) \sin t, \quad (11)$$

where A and B are arbitrary constants. Then (11) takes the following form

$$G_n(s) = \left(A - \int \left[\kappa_{n-2}(s)G_{n-2}(s) \sin \int \kappa_{n-1}(s)ds \right] ds \right) \cos \int \kappa_{n-1}(s)ds + \left(B + \int \left[\kappa_{n-2}(s)G_{n-2}(s) \cos \int \kappa_{n-1}(s)ds \right] ds \right) \sin \int \kappa_{n-1}(s)ds. \quad (12)$$

From (10), the function G_{n-1} is given by

$$G_{n-1}(s) = \left(A - \int \left[\kappa_{n-2}(s)G_{n-2}(s) \sin \int \kappa_{n-1}(s)ds \right] ds \right) \sin \int \kappa_{n-1}(s)ds - \left(B + \int \left[\kappa_{n-2}(s)G_{n-2}(s) \cos \int \kappa_{n-1}(s)ds \right] ds \right) \cos \int \kappa_{n-1}(s)ds. \quad (13)$$

From Equation (9) and (13), we have

$$\begin{aligned} \sum_{i=3}^{n-2} G_i G'_i &= G_3 \kappa_3 G_4 + G_4 \left(\kappa_4 G_5 - \kappa_3 G_3 \right) + \dots \\ &+ G_{n-3} \left(\kappa_{n-3} G_{n-2} - \kappa_{n-4} G_{n-4} \right) + G_{n-2} G'_{n-2} \\ &= G_{n-2} \left(G'_{n-2} + \kappa_{n-3} G_{n-3} \right) \\ &= \kappa_{n-2} G_{n-2} G_{n-1} \end{aligned}$$

If we integrate the above equation, we have

$$\sum_{i=3}^{n-2} G_i^2 = C - \left(A - \int \left[\kappa_{n-2}(s)G_{n-2}(s) \sin \int \kappa_{n-1}ds \right] ds \right)^2 - \left(B + \int \left[\kappa_{n-2}(s)G_{n-2}(s) \cos \int \kappa_{n-1}ds \right] ds \right)^2, \quad (14)$$

where C is a constant of integration. From Equations (12) and (13), we have

$$\begin{aligned} G_n^2 + G_{n-1}^2 &= \left(A - \int \left[\kappa_{n-2}(s) G_{n-2}(s) \sin \int \kappa_{n-1} ds \right] ds \right)^2 \\ &\quad + \left(B + \int \left[\kappa_{n-2}(s) G_{n-2}(s) \cos \int \kappa_{n-1} ds \right] ds \right)^2, \end{aligned} \quad (15)$$

It follows from (14) and (15) that

$$\sum_{i=3}^n G_i^2 = C.$$

Moreover this constant C calculates as follows. From (8), together the $(n-2)$ -equations (9), we have

$$C = \sum_{i=3}^n G_i^2 = \frac{1}{a_1^2} \sum_{i=3}^n a_i^2 = \frac{1 - a_1^2}{a_1^2} = \tan^2 \theta,$$

where we have used (2) and the fact that U is a unit vector field.

We do the converse of Theorem. Assume that the condition (9) is satisfied for a curve α . Let $\theta \in \mathbb{R}$ be so that $C = \tan^2 \theta$. Define the unit vector U by

$$U = \cos \theta \left[\mathbf{V}_1 + \sum_{i=3}^n G_i \mathbf{V}_i \right].$$

By taking account (9), a differentiation of U gives that $\frac{dU}{ds} = 0$, which it means that U is a constant vector field. On the other hand, the scalar product between the unit tangent vector field $\mathbf{V}_1$ with U is

$$\langle \mathbf{V}_1(s), U \rangle = \cos \theta.$$

Thus α is a cylindrical helix curve. This finishes the proof of Theorem 1.2.

As a direct consequence of the proof, we generalize Theorem 1.2 in Minkowski space and for timelike curves.

Theorem 2.1. *Let $\mathbf{E}_1^n$ be the Minkowski n -dimensional space and let $\alpha : I \rightarrow \mathbf{E}_1^n$ be a unit speed timelike curve. Then α is a cylindrical helix if and only if the function $\sum_{i=3}^n G_i^2$ is constant, where the functions G_i are defined as in (3).*

Proof. The proof carries the same steps as above and we omit the details. We only point out that the fact that α is timelike means that $\mathbf{V}_1(s) = \alpha'(s)$ is a timelike vector field. The other V_i in the Frenet frame, $2 \leq i \leq n$, are unit spacelike vectors and so, the second equation in (2) changes to $\mathbf{V}_2' = \kappa_1 \mathbf{V}_1 + \kappa_2 \mathbf{V}_3$ ([1, 6]).

□

3 Further characterizations of cylindrical helices

In this section we present new characterizations of cylindrical helix in $\mathbf{E}^n$. The first one is a consequence of Theorem 1.2.

Theorem 3.1. *Let $\alpha : I \subset \mathbb{R} \rightarrow \mathbf{E}^n$ be a unit speed curve in Euclidean space $\mathbf{E}^n$. Then α is a cylindrical helix if and only if there exists a C^2 -function $G_n(s)$ such that*

$$G_n = \frac{1}{\kappa_{n-1}} \left[\kappa_{n-2} G_{n-2} + G'_{n-1} \right], \quad \frac{dG_n}{ds} = -\kappa_{n-1}(s) G_{n-1}(s), \quad (16)$$

where

$$G_1 = 1, G_2 = 0, G_i = \frac{1}{\kappa_{i-1}} \left[\kappa_{i-2} G_{i-2} + G'_{i-1} \right], \quad 3 \leq i \leq n-1.$$

Proof. Let now assume that α is a cylindrical helix. By using Theorem 1.2 and by differentiation the (constant) function given in (4), we obtain

$$\begin{aligned} 0 &= \sum_{i=3}^n G_i G'_i \\ &= G_3 \kappa_3 G_4 + G_4 (\kappa_4 G_5 - \kappa_3 G_3) + \dots + G_{n-1} (\kappa_{n-1} G_n - \kappa_{n-2} G_{n-2}) + G_n G'_n \\ &= G_n (G'_n + \kappa_{n-1} G_{n-1}). \end{aligned}$$

This shows (16). Conversely, if (16) holds, we define a vector field U by

$$U = \cos \theta \left[\mathbf{V}_1 + \sum_{i=3}^n G_i \mathbf{V}_i \right].$$

By the Frenet equations (2), $\frac{dU}{ds} = 0$, and so, U is constant. On the other hand, $\langle \mathbf{V}_1(s), U \rangle = \cos \theta$ is constant, and this means that α is a cylindrical helix. $\square$

We end giving an integral characterization of a cylindrical helix.

Theorem 3.2. *Let $\alpha : I \subset \mathbb{R} \rightarrow \mathbf{E}^n$ be a unit speed curve in Euclidean space $\mathbf{E}^n$. Then α is a cylindrical helix if and only if the following condition is satisfied*

$$\begin{aligned} G_{n-1}(s) = & \left(A - \int \left[\kappa_{n-2} G_{n-2} \sin \int \kappa_{n-1} ds \right] ds \right) \sin \int^s \kappa_{n-1}(u) du \\ & - \left(B + \int \left[\kappa_{n-2} G_{n-2} \cos \int \kappa_{n-1} ds \right] ds \right) \cos \int^s \kappa_{n-1}(u) du. \end{aligned} \quad (17)$$

for some constants A and B .

Proof. Suppose that α is a cylindrical helix. By using Theorem 3.1, let define $m(s)$ and $n(s)$ by

$$\begin{aligned}\phi(s) &= \int^s \kappa_{n-1}(u) du, \\ m(s) &= G_n(s) \cos \phi + G_{n-1}(s) \sin \phi + \int \kappa_{n-2} G_{n-2} \sin \phi ds, \\ n(s) &= G_n(s) \sin \phi - G_{n-1}(s) \cos \phi - \int \kappa_{n-2} G_{n-2} \cos \phi ds.\end{aligned}\tag{18}$$

If we differentiate equations (18) with respect to s and taking into account of (17) and (16), we obtain $\frac{dm}{ds} = 0$ and $\frac{dn}{ds} = 0$. Therefore, there exist constants A and B such that $m(s) = A$ and $n(s) = B$. By substituting into (18) and solving the resulting equations for $G_{n-1}(s)$, we get

$$G_{n-1}(s) = \left(A - \int \kappa_{n-2} G_{n-2} \sin \phi ds \right) \sin \phi - \left(B + \int \kappa_{n-2} G_{n-2} \cos \phi ds \right) \cos \phi.$$

Conversely, suppose that (17) holds. In order to apply Theorem 3.1, we define $G_n(s)$ by

$$G_n(s) = \left(A - \int \kappa_{n-2} G_{n-2} \sin \phi ds \right) \cos \phi + \left(B + \int \kappa_{n-2} G_{n-2} \cos \phi ds \right) \sin \phi.$$

with $\phi(s) = \int^s \kappa_{n-1}(u) du$. A direct differentiation of (17) gives

$$G'_{n-1} = \kappa_{n-1} G_n - \kappa_{n-2} G_{n-2}.$$

This shows the left condition in (16). Moreover, a straightforward computation leads to $G'_n(s) = -\kappa_{n-1} G_{n-1}$, which finishes the proof. $\square$

We end this section with a characterization of cylindrical helices only in terms of the curvatures of α . From the definitions of G_i in (3), one can express the functions G_i in terms of G_3 and the curvatures of α as follows:

$$G_j = \sum_{i=0}^{j-3} A_{ji} G_3^{(i)}, \quad 3 \leq j \leq n,\tag{19}$$

where

$$G_3^{(i)} = \frac{d^{(i)} G_3}{ds^i}, \quad G_3^{(0)} = G_3 = \frac{\kappa_1}{\kappa_2}.$$

Then

$$\begin{aligned}G_4 &= \kappa_3^{-1} G'_3 = A_{41} G'_3 + A_{40} G_3, & A_{41} &= \kappa_3^{-1}, A_{40} = 0 \\ G_5 &= A_{52} G''_3 + A_{51} G'_3 + A_{50} G_3, & A_{52} &= \kappa_4^{-1} A_{41}, A_{51} = \kappa_4^{-1} A'_{41}, A_{50} = \kappa_4^{-1} \kappa_3\end{aligned}$$

and so on. Define the following functions:

$$A_{30} = 1, A_{40} = 0$$

$$A_{j0} = \kappa_{j-1}^{-1} \kappa_{j-2} A_{(j-2)0} + \kappa_{j-1}^{-1} A'_{(j-1)0}, \quad 5 \leq j \leq n$$

$$A_{j(j-3)} = \kappa_{j-1}^{-1} \kappa_{j-2}^{-1} \kappa_{j-3}^{-1} \dots \kappa_4^{-1} \kappa_3^{-1}, \quad 4 \leq j \leq n$$

$$A_{j(j-4)} = \kappa_{j-1}^{-1} \left(\kappa_{j-2}^{-1} \kappa_{j-3}^{-1} \dots \kappa_4^{-1} \kappa_3^{-1} \right)' + \kappa_{j-1}^{-1} \kappa_{j-2}^{-1} \left(\kappa_{j-3}^{-1} \dots \kappa_4^{-1} \kappa_3^{-1} \right)' \\ + \dots + \kappa_{j-1}^{-1} \kappa_{j-2}^{-1} \kappa_{j-3}^{-1} \dots \kappa_4^{-1} \left(\kappa_3^{-1} \right)', \quad 5 \leq j \leq n$$

$$A_{ji} = \kappa_{j-1}^{-1} \kappa_{j-2} A_{(j-2)i} + \kappa_{j-1}^{-1} \left(A'_{(j-1)i} + A_{(j-1)(i-1)} \right) \quad 1 \leq i \leq j-5, 6 \leq j \leq n$$

and $A_{ji} = 0$ otherwise.

The second equation of (16), leads the following condition:

$$A_{n(n-3)} G_3^{(n-2)} + \left(A'_{n(n-3)} + A_{n(n-4)} \right) G_3^{(n-3)} \\ + \sum_{i=1}^{n-4} \left[A'_{ni} + A_{n(i-1)} + \kappa_{n-1} A_{(n-1)i} \right] G_3^{(i)} \quad (20) \\ + \left(A'_{n0} + \kappa_{n-1} A_{(n-1)0} \right) G_3 = 0, \quad n \geq 3.$$

As a consequence of (20) and Theorem 1.2, we have the following

Corollary 3.3. *Let $\alpha : I \rightarrow \mathbf{E}^n$ be a unit speed curve in $\mathbf{E}^n$. The next statements are equivalent:*

1. α is a cylindrical helix.

2.

$$0 = A_{n(n-3)} \left(\frac{\kappa_1}{\kappa_2} \right)^{(n-2)} + \left(A'_{n(n-3)} + A_{n(n-4)} \right) \left(\frac{\kappa_1}{\kappa_2} \right)^{(n-3)} \\ + \sum_{i=1}^{n-4} \left[A'_{ni} + A_{n(i-1)} + \kappa_{n-1} A_{(n-1)i} \right] \left(\frac{\kappa_1}{\kappa_2} \right)^{(i)} \\ + \left(A'_{n0} + \kappa_{n-1} A_{(n-1)0} \right) \left(\frac{\kappa_1}{\kappa_2} \right), \quad n \geq 3.$$

3. The function

$$\sum_{j=3}^n \sum_{i=0}^{j-3} \sum_{k=0}^{j-3} A_{ji} A_{jk} \left(\frac{\kappa_1}{\kappa_2} \right)^{(i)} \left(\frac{\kappa_1}{\kappa_2} \right)^{(k)} = C$$

is constant, $j-i \geq 3, j-k \geq 3$.

References

- [1] A. Ali, R. López, Slant helices in Minkowski space $\mathbf{E}_1^3$, preprint 2008: arXiv:0810.1464v1 [math.DG].
- [2] H. Gluck, Higher curvatures of curves in Eulidean space, Amer. Math. Monthly, 73 (1996), 699–704.
- [3] A. Magden, On the curves of constant slope, YYÜ Fen Bilimleri Dergisi, 4 (1993), 103–109.
- [4] R. S. Milman, G. D. Parker, Elements of Differential Geometry, Prentice-Hall Inc., Englewood Cliffs, New Jersey, 1977.
- [5] E. Özdamar, H. H. Hacisalihoglu, A characterization of inclined curves in Euclidean n -space, Comm Fac Sci Univ Ankara, series A1, 24A (1975), 15–23.
- [6] M. Petrovic-Torgasev, E. Sucurovic, W-curves in Minkowski spacetime, Novi. Sad. J. Math. 32 (2002), 55–65.
- [7] P. D. Scofield, Curves of constant precession, Amer. Math. Monthly, 102 (1995), 531–537.